



---

## A NEW ITERATIVE APPROXIMATION OF A SPLIT FIXED POINT CONSTRAINT EQUILIBRIUM PROBLEM

MUSA ADEWALE OLANA<sup>1</sup>, ADHIR MAHARAJ<sup>2</sup> AND OJEN KUMAR NARAIN<sup>3</sup>

*Received 7 April, 2022; accepted 9 April, 2024; published 28 June, 2024.*

<sup>1</sup>SCHOOL OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE, UNIVERSITY OF KWAZULU-NATAL,  
DURBAN, SOUTH AFRICA,  
219095783@stu.ukzn.ac.za

<sup>2</sup>DEPARTMENT OF MATHEMATICS, DURBAN UNIVERSITY OF TECHNOLOGY, DURBAN, SOUTH AFRICA,  
adhirm@dut.ac.za

<sup>3</sup>SCHOOL OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE, UNIVERSITY OF KWAZULU-NATAL,  
DURBAN, SOUTH AFRICA,  
naraino@ukzn.ac.za

**ABSTRACT.** The purpose of this paper is to introduce an iterative algorithm for approximating an element in the solution set of the common split feasibility problem for fixed points of demimetric mappings and equilibrium problem for monotone mapping in real Hilbert spaces. Motivated by self-adaptive step size method, we incorporate the inertial technique to accelerate the convergence of the proposed method and establish a strong convergence of the sequence generated by the proposed algorithm. Finally, we present a numerical example to illustrate the significant performance of our method. Our results extend and improve some existing results in the literature.

*Key words and phrases:* Equilibrium problem; Demimetric mappings; Fixed point problems; Iterative method.

*2010 Mathematics Subject Classification.* 65K15, 47J25, 65J15, 90C33.

## 1. INTRODUCTION

Let  $C$  be a nonempty closed convex subset of a real Hilbert space  $H$  and  $S : C \rightarrow C$  be a nonlinear mapping. The Fixed Point Problem (FPP) is to find a point  $x \in C$  such that

$$(1.1) \quad Sx = x, \forall x \in C.$$

The fixed point set of the mapping  $S$  is denoted by  $Fix(S)$ . The fixed point theory finds its application in the prove of existence of solution of many nonlinear problems arising in many real life situations. From the existence of solution of differential equation to integral equations and evolutionary equations. The fixed point of many linear and nonlinear operators have been considered in the literature (see [16, 17, 18, 25]).

Let  $H_1$  and  $H_2$  be two real Hilbert spaces,  $S : H_1 \rightarrow H_1$  and  $T : H_2 \rightarrow H_2$  be nonlinear mappings. Let  $A : H_1 \rightarrow H_2$  be a bounded linear operator with its adjoint  $A^*$ , then, the following Split Common Fixed Point Problem (SCFPP) is to find:

$$(1.2) \quad \text{Finding } x \in Fix(S) \text{ such that } Ax \in Fix(T).$$

The SCFPP (1.2) initially introduced and studied by Censor and Segal [13], is a generalization of the Split Feasibility Problem (SFP) arising from signal processing and image restoration (see [12, 35]). Note that solving (1.2) can be translated to solving the following fixed point equation (see [1, 2, 3, 4, 5, 19]).

$$x^* = S(x^* - \tau A^*(I - T)Ax^*), \tau \geq 0.$$

Recently, Censor and Segal [13] proposed the following algorithm to solve SCFP (1.2):

**Algorithm 1.1.** : *Initialization:* Let  $x^* \in H_1 := \mathbb{R}_n$  be arbitrary. *Iterative step:* let

$$x_{n+1} = S(x_n - \tau A^*(I - T)Ax_n), n \geq 0$$

where  $S : \mathbb{R}_n \rightarrow \mathbb{R}_n$  and  $T : \mathbb{R}_m \rightarrow \mathbb{R}_m$  are two directed mappings and  $\tau \in (0, \frac{2}{\lambda})$  with  $\lambda$  being the spectral radius of the operator  $A^*A$ . There has been growing interest in the (SCFPP) due to its various applications, (see for example, [11, 36]).

In 2019, Chen *et al.* [14] introduced the following self-adaptive algorithm for solving SCFPP for demimetric mappings in real Hilbert spaces as follows

**Algorithm 1.2.** *Initialization:* Let  $x_0 \in H_1$  be arbitrary. For  $n \geq 0$ , assume the current iterate  $x_n$  has been constructed. If

$$\|x_n - Sx_n + A^*(I - T)Ax_n\| = 0,$$

then stop (in this case  $x_n$  solves problem (1.2)). Otherwise, calculate the next iterate  $x_{n+1}$  by the following formula

$$(1.3) \quad \begin{cases} y_n = x_n - Sx_n + A^*(I - T)Ax_n, \\ x_{n+1} = x_n - \alpha \tau_n y_n, \forall n \geq 0, \end{cases}$$

where  $\alpha \in (0, \min\{1 - \beta, 1 - \mu\})$  is a positive constant and  $\tau_n$  is chosen self adaptively as

$$\tau_n = \frac{\|x_n - Sx_n\|^2 + \|(I - T)Ax_n\|^2}{\|y_n\|^2}.$$

They assumed the sequence  $\{x_n\}$  generated by (1.3) is infinite and prove a weak convergence theorem for approximating the solution of the SCFPP.

In 1994, Blum and Oettli [10] introduced the notion of Equilibrium Problem (EP) as a generalization of certain optimization and variational inequality problems. It has received much attention from many researchers since its establishment due to its application to many problems arising from finance, physics, economics and so on. For this reason, several authors have introduced various generalizations of EP and numerous iterative algorithms have been developed, to solve these problems. The EP consists of finding a point  $x \in C$  such

$$(1.4) \quad F(x, y) \geq 0, \quad \forall y \in C,$$

where  $F : C \times C \rightarrow \mathbb{R}$  is a bifunction.

The Mixed Equilibrium Problem (MEP) which is a generalization of the EP (1.3) is known to include fixed point problem, optimization problem, variational inequality problem, and Nash equilibrium problem as special cases; (see [10, 20]). Some methods have been proposed to solve the MEP, see, for example, [20, 22]. The Mixed Equilibrium Problem (MEP) is to find  $x^* \in C$  such that

$$(1.5) \quad F(x^*, x) + \phi(x) - \phi(x^*) \geq 0 \text{ for all } x \in C.$$

If in (1.5)  $\phi = 0$ , then the MEP (1.5) reduces to the EP (1.4).

In 2017, Wang [34] introduced the following new iterative algorithm for the SCFPP of directed mappings

**Algorithm 1.3.** choose an arbitrary initial guess  $x_0$ . Assume  $x_n$  has been constructed. If

$$\|x_n - Sx_n + A^*(I - T)Ax_n\| = 0,$$

then stop; otherwise, continue and construct  $x_{n+1}$  via the formula;

$$x_{n+1} = x_n - \tau_n \{\|x_n - Sx_n + A^*(I - T)Ax_n\|\}, \quad \forall n \geq 0,$$

where  $\tau_n$  is choose self-adaptively as

$$\tau_n = \frac{\|x_n - Sx_n\|^2 + \|(I - T)Ax_n\|^2}{\|x_n - Sx_n + A^*(I - T)Ax_n\|^2}$$

**Algorithm 1.4.** Let  $u \in H$  and start an initial guess  $x_0 \in H$ , assume  $x_n$  has been constructed. If

$$\|x_n - Sx_n + A^*(I - T)Ax_n\| = 0,$$

then stop; otherwise, continue and construct  $x_{n+1}$  via the formula;

$$x_{n+1} = \alpha_n u + (1 - \alpha_n) \{x_n - Sx_n + A^*(I - T)Ax_n\}, \quad \forall n \geq 0,$$

where stepsize sequence  $\tau_n$  is choose self-adaptively as

$$\tau_n = \frac{\|x_n - Sx_n\|^2 + \|(I - T)Ax_n\|^2}{\|x_n - Sx_n + A^*(I - T)Ax_n\|^2}$$

Wang obtained a weak and a strong convergence of Algorithms 1.3 and 1.4, respectively. Wang's results in [34] from the directed mappings to the demicontractive mappings. Further, they construct the following two self-adaptive algorithms for solving the split common fixed point problem (1.2).

**Algorithm 1.5.** . Initialization: Let  $x_0 \in H_1$  be arbitrary . For  $n \geq 0$  , assume the current iterate  $x_n$  has been constructed. If

$$\|x_n - Sx_n + A^*(I - T)Ax_n\| = 0,$$

then stop; otherwise , calculate the next iterate  $x_{n+1}$  by the following formula

$$\begin{cases} y_n = x_n - Sx_n + A^*(I - T)Ax_n, \\ z_n = x_n - \alpha_n \tau_n y_n, \forall n \geq 0, \end{cases}$$

where  $\alpha \in (0, \min\{1 - \beta, 1 - \mu\})$  is a positive constant and  $\tau_n$  is chosen self-adaptively as

$$\tau_n = \frac{\|x_n - Sx_n\|^2 + \|(I - T)Ax_n\|^2}{\|y_n\|^2}$$

**Algorithm 1.6.** Initialization: let  $u \in H_1$  be a fixed point and let  $x_0 \in H$  be arbitrary. Iterative step: for  $n \geq 0$  , assume the current iterate  $x_n$  has been constructed . If

$$\|x_n - Sx_n + A^*(I - T)Ax_n\| = 0,$$

then stop; otherwise , calculate the next iterate  $x_{n+1}$  by the following formula

$$\begin{cases} y_n = x_n - Sx_n + A^*(I - T)Ax_n, \\ x_{n+1} = \alpha_n u + (1 - \alpha_n)(x_n - \alpha \tau_n y_n), \forall n \geq 0, \end{cases}$$

where  $\alpha \in (0, \min\{1 - \beta, 1 - \mu\})$  is a positive constant and  $\tau_n$  is chosen self-adaptively as

$$\tau_n = \frac{\|x_n - Sx_n\|^2 + \|(I - T)Ax_n\|^2}{\|y_n\|^2}$$

They also obtained a weak and a strong convergence result of Algorithms 1.5 and 1.6, respectively. They present two self-adaptive algorithms for solving the split common fixed point problem (1.2).

Recently, Shehu introduced a hybrid method for finding a common fixed point of infinite family of  $k$ -strictly pseudocontractive mappings, the set of common solutions to a system of generalized mixed equilibrium problem, and the set of solutions to variational inequality problem in Hilbert space. Starting with an arbitrary  $x_0 \in C$ ,  $C_{1,i} = C$ ,  $C_1 = \bigcap_{i=1}^{\infty} C_{1,i}$ , and  $x_1 = P_{C_1} x_0$  define sequence  $\{x_n\}$ ,  $\{w_n\}$ ,  $\{u_n\}$ ,  $\{z_n\}$ , and  $\{y_{n,i}\}$  as follows:

$$(1.6) \quad \begin{cases} z_n & = T_{r_n}^{F_1, \phi_1}(x_n - r_n Ax_n), \\ y_n & = T_{r_n}^{F_2, \phi_2}(z_n - \lambda_n Bx_n), \\ w_n & = P_C(u_n - s_n D u_n), \\ y_{n,i} & = \alpha_{n_i} w_n + (1 - \alpha_{n_i}) T_i w_n, \quad n \geq 1, \\ C_{n+1,i} & = \{z \in C_{n,i} : \|y_{n,i} - z\| \leq \|x_n - z\|\}, \quad n \geq 1, \\ C_{n+1} & = \bigcap_{i=1}^{\infty} C_{n+1,i}, \\ x_{n+1} & = P_{C_{n+1}} x_0, \quad n \geq 1, \end{cases}$$

where  $T_i$  is a  $k_i$ -strictly pseudocontractive mapping and for some  $0 \leq k_i < 1$ ,  $A, B$  is  $\alpha, \beta$ -inverse- strongly monotone mapping of  $C$  into  $H$ . He proved that if the sequence  $\{\alpha_{n_i}\}$ ,  $\{r_n\}$ ,  $\{s_n\}$  and  $\{\lambda_n\}$  of parameters satisfies appropriate conditions, then  $\{x_n\}$  generated by (1.6)

Moudafi [24] recently studied the convergence properties of a relaxed algorithm for SCFP for a class of quasi-nonexpansive operators  $T$  such that  $I - T$  is demiclosed at zero. He also proved a weak convergence theorem as shown below.

**Theorem 1.1.** *Given a bounded linear operator  $A : H_1 \rightarrow H_2$ , let  $U : H_1 \rightarrow H_1$  and  $T : H_2 \rightarrow H_2$  be two quasi-nonexpansive operators with nonempty sets  $\text{Fix}(U) = C$  and  $\text{Fix}(T) = Q$ . Assume that  $I - U$  and  $I - T$  are demiclosed at zero. Suppose  $\Gamma := \{x \in C : Ax \in Q\} \neq \emptyset$  and define an iterative sequence  $x_n$  by*

$$(1.7) \quad \begin{cases} x_0 \in H_1, \\ u_n = x_n + \alpha\beta A^*(T - I)Ax_n, \\ x_{n+1} = (1 - \alpha_n)u_n + \alpha_n U(u_n). \end{cases}$$

Polyak [26] first proposed an inertial extrapolation as an acceleration process to solve the smooth convex minimization problem. The inertial algorithm is a two-step iteration where the next iterate is defined by making use of the previous two iterates. Recently, several researchers have constructed some fast iterative algorithms by using inertial extrapolation (see, e.g., [6, 7])

Motivated by the above results and the current research interest in this direction, in this article, we propose a new iterative scheme for approximating an element in the solution set of the common split feasibility problem for fixed points of demimetric mappings and equilibrium problem for monotone mapping in a real Hilbert space. We incorporate self-adaptive step size method and inertial technique to accelerate the convergence of the proposed method, we establish the strong convergence of the sequence generated by the proposed algorithm. We finally, establish some applications and numerical examples to illustrate the significant performance of our method.

Subsequent sections of this work are organized as follows: In Section 2, we recall some basic definitions and Lemmas that are relevant in establishing our main results. In Section 3, we state some Lemmas that are useful in establishing the strong convergence of our proposed algorithm and also prove the strong convergence theorem for the algorithm.

## 2. PRELIMINARIES

In this paper, let  $H$  be a real Hilbert space with inner product  $\langle \cdot, \cdot \rangle$  and the norm given by  $\|\cdot\|$  respectively. We denote the weak and strong convergence of a sequence  $x_n$  to a point  $x \in H$  by  $x_n \rightharpoonup x$  and  $x_n \rightarrow x$  respectively.

**Definition 2.1.** A mapping  $A : H \rightarrow H$  is said to be:

(i) monotone if:

$$\langle Ax - Ay, x - y \rangle \geq 0, \quad \forall x, y \in H;$$

(ii)  $\lambda$ -inverse strongly monotone (co-coercive) if there exists  $\lambda > 0$  such that:

$$\langle Ax - Ay, x - y \rangle \geq \lambda \|Ax - Ay\|^2, \quad \forall x, y \in H;$$

(iii) nonexpansive if:

$$\|Ax - Ay\| \leq \|x - y\|, \quad \forall x, y \in H;$$

(iv) firmly nonexpansive if:

$$\langle Ax - Ay, x - y \rangle \geq \|Ax - Ay\|^2, \quad \forall x, y \in H.$$

For  $x \in H$ , there exists the unique nearest point  $P_C x$  in  $C$  such that

$$\|x - y\| \leq \|x - P_C x\|, \quad \forall y \in C.$$

$P_C$  is called metric projection of  $H$  onto  $C$ . It is known that  $P_C$  is nonexpansive.

**Lemma 2.1.** [15] *Let  $C$  be a closed and convex subset in a real Hilbert space  $H$ , for any  $x \in H$  and  $z \in C$ , we have*

$$z = P_C x \Leftrightarrow \langle x - z, z - y \rangle \geq 0, \forall y \in C.$$

**Lemma 2.2.** [15] *Let  $C$  be a closed and convex subset in a real Hilbert space  $H$  and let  $x \in H$ , then we have the following*

- (i)  $\|P_C x - P_C y\|^2 \leq \langle P_C x - P_C y, x - y \rangle \forall y \in H.$
- (ii)  $\|P_C x - y\|^2 \leq \|x - y\|^2 - \|x - P_C x\|^2 \forall y \in C.$

For more properties of the metric projection, refer [15, Section 3].

We need the following assumptions to solve a mixed equilibrium problem for a bifunction  $F : C \times C \rightarrow \mathbb{R}$  and a mapping  $\phi : C \rightarrow \mathbb{R}$ .

- (A1)  $F(x, x) = 0, \forall x \in C,$
- (A2)  $F$  is monotone, i.e.,  $F(x, y) + F(y, x) \leq 0 \forall x, y \in C,$
- (A3)  $\lim_{\alpha \rightarrow \infty} F(\alpha z + (1 - \alpha)x, y) \leq F(x, y) \forall x, y, z \in C,$
- (A4)  $\forall x \in C, y \mapsto F_1(x, y)$  is convex and lower semicontinuous,
- (A5) for each  $x \in C, \alpha \in (0, 1]$ , and  $r > 0$ , there exist a bounded subset  $D \subseteq C$  and  $y \in C$  such that for any  $z \in C/D$ ,

$$F(z, y) + \phi(y) - \phi(z) + \frac{1}{r} \langle y - z, z - x \rangle < 0.$$

- (A6)  $C$  is a bounded set.

**Lemma 2.3.** [28] *Let  $C$  be a nonempty closed convex subset of a Hilbert space  $H_1$  and  $\phi : C \rightarrow \mathbb{R} \cup \{+\infty\}$  a proper lower semicontinuous and convex mapping such that  $C \cap \text{dom } \phi = \emptyset$ . Suppose that bifunction  $F : C \times C \rightarrow \mathbb{R}$  and a mapping  $\phi$  satisfy Conditions (A1)(A6). For  $r > 0$  and  $x \in H$ , let  $T_r^{F, \phi} : H \rightarrow C$  be a mapping defined by*

$$(2.1) \quad T_r^{F, \phi}(x) = \{z \in C : F(z, y) + \phi(y) - \phi(z) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \forall y \in C\}.$$

Assume that either (A5) or (A6) holds. Then:

- (i) for each  $x \in H, T_r^{F, \phi} x \neq \emptyset,$
- (ii)  $T_r^{F, \phi}$  is single valued,
- (iii)  $T_r^{F, \phi}$  is firmly nonexpansive,
- (iv)  $\text{Fix}(T_r^{F, \phi}) = \text{MEP}(F, \phi)$  and it is closed and convex.

**Lemma 2.4.** [21] *Let  $H$  be a real Hilbert space. Then, we have*

$$\begin{aligned} \|x - y\|^2 &= \|x\|^2 - \|y\|^2 - 2\langle x - y, y \rangle, \\ \|x + y\|^2 &\leq \|x\|^2 + 2\langle y, x + y \rangle, \end{aligned}$$

and

$$\|\alpha x + (1 - \alpha)y\|^2 = \alpha\|x\|^2 + (1 - \alpha)\|y\|^2 - \alpha(1 - \alpha)\|x - y\|^2$$

for all  $x, y \in H$  and  $\alpha \in [0, 1]$ . Also, if  $\{x_n\}$  is a sequence in  $H$  weakly converging to  $z \in H$ , then

$$\limsup_{n \rightarrow \infty} \|x_n - y\|^2 = \limsup_{n \rightarrow \infty} \|x_n - z\|^2 + \|z - y\|^2, \forall y \in H.$$

**Lemma 2.5.** [29] *Let  $\{a_n\} \subset \mathbb{R}_+, \{b_n\} \subset \mathbb{R}$  and  $\{\xi_n\} \subset (0, 1)$  be such that  $\sum_{n=1}^{\infty} \xi_n = \infty$  and*

$$a_{n+1} \leq (1 - \xi_n)a_n + \xi_n b_n, \forall n \in \mathbb{N}.$$

If  $\limsup_{i \rightarrow \infty} b_{n_i} \leq 0$  for every subsequence  $a_{n_i}$  of  $a_n$  satisfying  $\liminf_{i \rightarrow \infty} (a_{n_{i+1}} - a_{n_i}) \geq 0$ , then  $\lim_{n \rightarrow \infty} a_n = 0$ .

### 3. MAIN RESULT

We study the problem of finding a common solution of the split common fixed points problem and mixed equilibrium problem(SCFPPMEP). We denote by  $\Gamma$  the solution set of the SCFPPMEP. That is  $\Gamma := \{x \in C : x \in \text{Fix}(S) \cap \text{MEP}(F, \phi) \text{ such that } Ax \in \text{Fix}(T)\}$ . For solving the SCFPPMEP, we make the following assumptions:

- (B1) Let  $C$  and  $Q$  be nonempty, closed and convex subsets of real Hilbert spaces  $H_1$  and  $H_2$  respectively;
- (B2)  $S : C \rightarrow C$  and  $T : Q \rightarrow Q$  are two demimetric mappings with constants  $\beta \in (-\infty, 1)$  and  $\mu \in (-\infty, 1)$ , respectively;
- (B3)  $A : H_1 \rightarrow H_2$  is bounded linear operator with its adjoint operator  $A^*$ ;
- (B4) The bifunction  $F : C \times C \rightarrow \mathbb{R}$  and  $\phi : C \rightarrow \mathbb{R}$  satisfy condition (A1) - (A6);
- (B5)  $\Gamma$  is nonempty.

Next we proof the following self adaptive algorithm for solving the SCFPPMEP:

---

**Algorithm 1.** *Inertial Algorithm for SCFPPMEP*

---

**Initialization:** Choose  $x_0, x_1 \in C$ ,  $\theta \in (0, 1)$ ,  $\beta, \mu \in (0, 1)$ ,  $\{r_n\}$  a sequence of nonnegative real numbers and  $\alpha_n, \beta_n \subset (0, 1)$  satisfying

- (C1)  $\lim_{n \rightarrow \infty} \alpha_n = 0$  and  $\sum_{n=1}^{\infty} \alpha_n = \infty$ ;
- (C2)  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$ ;
- (C3)  $0 < \liminf_{n \rightarrow \infty} r_n$ ;
- (C4)  $\lim_{n \rightarrow \infty} \frac{\epsilon_n}{\alpha_n} = 0$ , where  $\epsilon_n$  is a sequence of nonnegative real numbers.

**Step 1:** Compute the inertial step

$$(3.1) \quad \bar{\theta}_n = \begin{cases} \min\{\theta, \frac{\epsilon_n}{\|x_n - x_{n-1}\|}\}, & \text{if } x_n \neq x_{n-1} \\ \theta, & \text{otherwise.} \end{cases}$$

$$w_n = x_n + \bar{\theta}_n(x_n - x_{n-1}).$$

**Step 2:** Compute

$$(3.2) \quad \begin{cases} y_n = \beta_n w_n + (1 - \beta_n)T_{r_n}^{F, \phi} w_n, \\ z_n = y_n - S y_n + A^*(A y_n - T A y_n), \\ x_{n+1} = \alpha_n f(x_n) + (1 - \alpha_n)(y_n - \alpha \tau_n z_n), \end{cases}$$

where  $\alpha \in (0, \min\{1 - \beta, 1 - \mu\})$  is a positive constant and  $\tau_n$  is chosen self-adaptively as

$$\tau_n = \frac{(\|y_n - S y_n\|^2 + \|A y_n - T A y_n\|^2)}{\|z_n\|^2}.$$

**Remark 1.** We assume that the sequence  $\{x_n\}$  generated by Algorithm 1 is infinite. In other words, Algorithm 1 does not terminate in a finite number of iterations.

**Lemma 3.1.** If  $\|y_n - S y_n + A^*(I - T)A y_n\| = 0$ , then we arrive at the solution of the SCFPP.

*Proof:* If  $y_n$  solves split common fixed points problem, then  $y_n = Sy_n$  and  $(I - T)Ay_n = 0$ , Therefore, we have  $\|y_n - Sy_n + A^*(I - T)Ay_n\| = 0$ . To see the converse, suppose that  $\|y_n - Sy_n + A^*(I - T)Ay_n\| = 0$ . Then we have  $y \in \Omega$  such that

$$\begin{aligned} 0 &= \|y_n - Sy_n + A^*(I - T)Ay_n\| \|y_n - y\| \\ &\geq \langle y_n - Sy_n + A^*(I - T)Ay_n, y_n - y \rangle \\ &\geq \langle y_n - Sy_n, y_n - y \rangle + \langle A^*(I - T)Ay_n, y_n - y \rangle \\ (3.3) \quad &\geq \langle y_n - Sy_n, y_n - y \rangle + \langle (I - T)Ay_n, Ay_n - Ay \rangle. \end{aligned}$$

Since  $S$  and  $T$  are demimetric, we have that

$$(3.4) \quad \langle y_n - Sy_n, y_n - y \rangle \geq \frac{1 - \beta}{2} \|y_n - Sy_n\|^2$$

and

$$(3.5) \quad \langle (I - T)Ay_n, Ay_n - Ay \rangle \geq \frac{1 - \mu}{2} \|Ay_n - TAy_n\|^2$$

we obtain the following by combining (3.3), (3.4) and (3.5),

$$(3.6) \quad 0 \geq \frac{1 - \beta}{2} \|y_n - Sy_n\|^2 + \frac{1 - \mu}{2} \|Ay_n - TAy_n\|^2$$

Since  $\beta, \mu \in (-\infty, 1)$ , we infer that  $y_n \in \text{Fix}(S)$  and  $Ay_n \in \text{Fix}(T)$  by (3.6). Therefore,  $y_n$  solves problem of common fixed point problem. This completes the proof.

**Theorem 3.2.** *Suppose assumption (B1)-(B5) hold. Then the sequence  $\{x_n\}$  generated by Algorithm 1 converges strongly to an element  $p = P_\Gamma f(p) \in \Gamma$ .*

*Proof:* Fix  $p \in \Omega$ , then from (1), we have

$$\begin{aligned} \langle z_n, y_n - p \rangle &= \langle y_n - Sy_n + A^*(Ay_n - TAy_n), y_n - p \rangle \\ &= \langle y_n - Sy_n, y_n - p \rangle + \langle A^*(Ay_n - TAy_n), y_n - p \rangle \\ &\geq \frac{1 - \beta}{2} \|y_n - Sy_n\|^2 + \frac{1 - \mu}{2} \|Ay_n - TAy_n\|^2 \\ &\geq \frac{1}{2} \min\{1 - \beta, 1 - \mu\} (\|y_n - Sy_n\|^2 + \|Ay_n - TAy_n\|^2). \end{aligned}$$

Now let  $v_n = y_n - \alpha\tau_n z_n - p$ , we get

$$\begin{aligned} \|v_n - p\|^2 &= \|y_n - \alpha\tau_n z_n - p\|^2 \\ &= \|y_n - p\|^2 - 2\alpha\tau_n \langle z_n, y_n - p \rangle + \alpha^2\tau_n^2 \|y_n\|^2 \\ &= \|y_n - p\|^2 + \alpha^2 \frac{(\|y_n - Sy_n\|^2 + \|Ay_n - TAy_n\|^2)}{\|z_n\|^2} \\ &\quad - \alpha \min\{1 - \beta, 1 - \mu\} \frac{(\|y_n - Sy_n\|^2 + \|Ay_n - TAy_n\|^2)}{\|z_n\|^2} \\ (3.7) \quad &\leq \|y_n - p\|^2 - \alpha \min\{1 - \beta, 1 - \mu\} \frac{(\|y_n - Sy_n\|^2 + \|Ay_n - TAy_n\|^2)}{\|z_n\|^2} \\ &\leq \|y_n - p\|^2. \end{aligned}$$

Thus, we obtain that

$$\|v_n - p\| \leq \|y_n - p\|.$$



Observe that

$$\begin{aligned} \|y_n - p\| &= \|\beta_n w_n + (1 - \beta_n)T_{r_n}^{F,\phi} w_n - p\| \\ &\leq \beta_n \|w_n - p\| + (1 - \beta_n) \|T_{r_n}^{F,\phi} w_n - p\| \\ &\leq \beta_n \|w_n - p\| + (1 - \beta_n) \|w_n - p\| \\ &= \|w_n - p\|. \end{aligned}$$

Furthermore, we have

$$\begin{aligned} \|x_{n+1} - p\| &= \|\alpha_n f(x_n) + (1 - \alpha_n)v_n - p\| \\ &\leq \alpha_n \|f(x_n) - f(p)\| + \alpha_n \|f(p) - p\| + (1 - \alpha_n) \|v_n - p\| \\ &\leq \alpha_n c \|x_n - p\| + \alpha_n \|f(p) - p\| + (1 - \alpha_n) \|v_n - p\| \\ &\leq \{1 - \alpha_n(1 - c)\} \|x_n - p\| + \alpha_n \|f(p) - p\| + \theta_n(1 - \alpha_n) \|x_n - x_{n-1}\| \\ &\leq \{1 - \alpha_n(1 - c)\} \|x_n - p\| + \alpha_n \left( \frac{\theta_n}{\alpha_n} \|x_n - x_{n-1}\| + \|f(p) - p\| \right) \\ &\leq \max \left\{ \|x_n - p\|, \frac{\frac{\theta_n}{\alpha_n} \|x_n - x_{n-1}\| + \|f(p) - p\|}{1 - c} \right\}. \end{aligned}$$

Therefore, we obtain by (3.1) and (C4), that  $\frac{\theta_n}{\alpha_n} \|x_n - x_{n-1}\| \rightarrow 0$  as  $n \rightarrow \infty$  and there exist  $M > 0$  such that  $\frac{\theta_n}{\alpha_n} \|x_n - x_{n-1}\| \leq M$  for all  $n \in \mathbb{N}$ . Hence,

$$\begin{aligned} \|x_{n+1} - p\| &\leq \max \left\{ \|x_n - p\|, \frac{M + \|f(p) - p\|}{1 - c} \right\} \\ &\leq \max \left\{ \|x_n - p\|, \frac{M + \|f(p) - p\|}{1 - c} \right\} \quad \forall n \in \mathbb{N}. \end{aligned}$$

Hence,  $\{x_n\}$  is bounded. It is easy to see that operator  $P_\Gamma f$  is a contraction. Thus by the Banach contraction principle, there exists a unique point  $p = P_\Gamma f(p)$ . It follows from the characterization of  $P_\Gamma$  that

$$(3.8) \quad \langle f(p) - p, q - p \rangle \leq 0, \quad \forall q \in \Gamma.$$

Using Lemma 2.4 and (3.7)

$$\begin{aligned}
\|x_{n+1} - p\|^2 &\leq \|(1 - \alpha_n)(v_n - p) + \alpha_n(f(x_n) - f(p))\|^2 + 2\alpha_n\langle f(p) - p, x_{n+1} - p \rangle \\
&\leq (1 - \alpha_n)\|v_n - p\|^2 + \alpha_n\|f(x_n) - f(p)\|^2 + 2\alpha_n\langle f(p) - p, x_{n+1} - p \rangle \\
&\leq (1 - \alpha_n)\|y_n - p\|^2 - \alpha(\min\{1 - \beta, 1 - \mu\}) \frac{(\|y_n - Sy_n\|^2 + \|Ay_n - TAy_n\|^2)}{\|z_n\|^2} \\
&\quad + \alpha_n\|f(x_n) - f(p)\|^2 + 2\alpha_n\langle f(p) - p, x_{n+1} - p \rangle \\
&\leq (1 - \alpha_n)\|w_n - p\|^2 - \alpha(\min\{1 - \beta, 1 - \mu\}) \frac{(\|y_n - Sy_n\|^2 + \|Ay_n - TAy_n\|^2)}{\|z_n\|^2} \\
&\quad + \alpha_n\|f(x_n) - f(p)\|^2 + 2\alpha_n\langle f(p) - p, x_{n+1} - p \rangle \\
&\leq (1 - \alpha_n)\|x_n - p\|^2 + 2\theta_n\langle x_n - p, x_n - x_{n-1} \rangle + \theta_n\|x_n - x_{n-1}\|^2 \\
&\quad - \alpha(\min\{1 - \beta, 1 - \mu\}) \frac{(\|y_n - Sy_n\|^2 + \|Ay_n - TAy_n\|^2)}{\|z_n\|^2} \\
&\quad + \alpha_n\|f(x_n) - f(p)\|^2 + 2\alpha_n\langle f(p) - p, x_{n+1} - p \rangle \\
&\leq (1 - \alpha_n)\|x_n - p\|^2 + \alpha_n(1 - c)b_n \\
(3.9) \quad &- \alpha(\min\{1 - \beta, 1 - \mu\}) \frac{(\|y_n - Sy_n\|^2 + \|Ay_n - TAy_n\|^2)}{\|z_n\|^2},
\end{aligned}$$

where

$$b_n := \frac{1}{1 - c} (2\langle f(p) - p, x_{n+1} - p \rangle + \frac{\theta_n^2}{\alpha_n} \|x_n - x_{n-1}\|^2 + 2\frac{\theta_n}{\alpha_n} \|x_n - p\| \|x_n - x_{n-1}\|).$$

It follows that

$$(3.10) \quad \alpha(\min\{1 - \beta, 1 - \mu\}) \frac{(\|y_n - Sy_n\|^2 + \|Ay_n - TAy_n\|^2)}{\|z_n\|^2} \leq \|x_n - p\|^2 - \|x_{n+1} - p\|^2 + \alpha_n(1 - c)M',$$

where  $M' = \sup\{b_n : n \in \mathbb{N}\}$ .

Now, set  $a_n = \|x_n - p\|^2$  and  $\eta_n := \alpha_n(1 - c)$ . From (3.9) we have the following inequality:

$$a_{n+1} \leq (1 - \eta_n)a_n + \eta_n b_n.$$

To apply Lemma 2.5, we have to show that  $\limsup_{i \rightarrow \infty} b_{n_i} \leq 0$  for every subsequence  $\{a_{n_i}\}$  of  $\{a_n\}$  satisfying

$$(3.11) \quad \liminf_{i \rightarrow \infty} (a_{n_{i+1}} - a_{n_i}) \geq 0.$$

To do this, suppose that  $\{a_{n_i}\} \subseteq \{a_n\}$  is a subsequence satisfying (3.11). Therefore, by (3.10) and (Cii), we have

$$\begin{aligned}
&\limsup_{i \rightarrow \infty} \alpha(\min\{1 - \beta, 1 - \mu\}) \frac{(\|y_{n_i} - Sy_{n_i}\|^2 + \|Ay_{n_i} - TAy_{n_i}\|^2)}{\|z_{n_i}\|^2} \\
&\leq \limsup_{i \rightarrow \infty} (a_{n_{i+1}} - a_{n_i}) + (1 - c)M' \lim_{i \rightarrow \infty} \alpha_{n_i} \\
&= -\liminf_{i \rightarrow \infty} (a_{n_{i+1}} - a_{n_i}) \\
&\leq 0,
\end{aligned}$$

which implies

$$(3.12) \quad \frac{(\|y_{n_i} - Sy_{n_i}\|^2 + \|Ay_{n_i} - TAy_{n_i}\|^2)}{\|z_{n_i}\|^2} = 0.$$

Taking into consideration that

$$(3.13) \quad \frac{(\|y_{n_i} - Sy_{n_i}\|^2 + \|Ay_{n_i} - TAy_{n_i}\|^2)}{2 \max\{1, \|A\|^2\}} \leq \frac{(\|y_{n_i} - Sy_{n_i}\|^2 + \|Ay_{n_i} - TAy_{n_i}\|^2)}{\|z_{n_i}\|^2}.$$

We deduce from (3.12), that

$$(3.14) \quad \lim_{i \rightarrow \infty} \|y_{n_i} - Sy_{n_i}\|^2 = \lim_{i \rightarrow \infty} \|Ay_{n_i} - TAy_{n_i}\|^2 = 0.$$

Observe from (3.2) and the nonexpansive property of  $T_{r_n}^{F,\phi}$ , that

$$(3.15) \quad \begin{aligned} \|y_n - p\|^2 &= \|\beta_n w_n + (1 - \beta_n)T_{r_n}^{F,\phi} w_n - p\|^2 \\ &= \beta_n \|w_n - p\|^2 + (1 - \beta_n) \|T_{r_n}^{F,\phi} w_n - p\|^2 - \beta_n(1 - \beta_n) \|w_n - T_{r_n}^{F,\phi} w_n\|^2 \\ &\leq \beta_n \|w_n - p\|^2 + (1 - \beta_n) \|w_n - p\|^2 - \beta_n(1 - \beta_n) \|w_n - T_{r_n}^{F,\phi} w_n\|^2 \\ &= \|w_n - p\|^2 - \beta_n(1 - \beta_n) \|w_n - T_{r_n}^{F,\phi} w_n\|^2. \end{aligned}$$

Again, by using Lemma 2.4, (3.7) and (3.15), we have that

$$(3.16) \quad \begin{aligned} \|x_{n+1} - p\|^2 &\leq \|(1 - \alpha_n)(v_n - p) + \alpha_n(f(x_n) - f(p))\|^2 + 2\alpha_n \langle f(p) - p, x_{n+1} - p \rangle \\ &\leq (1 - \alpha_n) \|v_n - p\|^2 + \alpha_n \|f(x_n) - f(p)\|^2 + 2\alpha_n \langle f(p) - p, x_{n+1} - p \rangle \\ &\leq (1 - \alpha_n) \|y_n - p\|^2 + \alpha_n \|f(x_n) - f(p)\|^2 + 2\alpha_n \langle f(p) - p, x_{n+1} - p \rangle \\ &\leq (1 - \alpha_n) (\|w_n - p\|^2 - \beta_n(1 - \beta_n) \|w_n - T_{r_n}^{F,\phi} w_n\|^2) + \alpha_n \|f(x_n) - f(p)\|^2 \\ &\quad + 2\alpha_n \langle f(p) - p, x_{n+1} - p \rangle \\ &\leq (1 - \alpha_n) \|x_n - p\|^2 + 2\theta_n \langle x_n - p, x_n - x_{n-1} \rangle + \theta_n \|x_n - x_{n-1}\|^2 \\ &\quad - \beta_n(1 - \beta_n) \|w_n - T_{r_n}^{F,\phi} w_n\|^2 + \alpha_n \|f(x_n) - f(p)\|^2 + 2\alpha_n \langle f(p) - p, x_{n+1} - p \rangle \\ &\leq [1 - \alpha_n(1 - c)] \|x_n - p\|^2 + \alpha_n(1 - c) b_n - \beta_n(1 - \beta_n) \|w_n - T_{r_n}^{F,\phi} w_n\|^2, \end{aligned}$$

where

$$b_n := \frac{1}{1 - c} (2 \langle f(p) - p, x_{n+1} - p \rangle + \frac{\theta_n^2}{\alpha_n} \|x_n - x_{n-1}\|^2 + 2 \frac{\theta_n}{\alpha_n} \|x_n - p\| \|x_n - x_{n-1}\|).$$

Thus, we obtain

$$(3.17) \quad \beta_n(1 - \beta_n) \|w_n - T_{r_n}^{F,\phi} w_n\|^2 \leq \|x_n - p\|^2 - \|x_{n+1} - p\|^2 + \alpha_n(1 - c) M',$$

where  $M' = \sup\{b_n : n \in \mathbb{N}\}$ .

Again as before, let  $a_n = \|x_n - p\|^2$  and  $\eta_n := \alpha_n(1 - c)$ . From (3.16), we have that

$$a_{n+1} \leq (1 - \eta_n) a_n + \eta_n b_n.$$

Therefore, by (C1) and (C2), we have

$$\begin{aligned} &\limsup_{i \rightarrow \infty} \beta_{n_i} (1 - \beta_{n_i}) \|w_{n_i} - T_{r_{n_i}}^{F,\phi} w_{n_i}\|^2 \\ &\leq \limsup_{i \rightarrow \infty} (a_{n_i+1} - a_{n_i}) + (1 - c) M' \lim_{i \rightarrow \infty} \alpha_{n_i} \\ &= - \liminf_{i \rightarrow \infty} (a_{n_i+1} - a_{n_i}) \\ &\leq 0, \end{aligned}$$

which implies

$$(3.18) \quad \lim_{i \rightarrow \infty} \|w_{n_i} - T_{r_{n_i}}^{F,\phi} w_{n_i}\| = 0.$$

Now,

$$\begin{aligned} \|w_n - x_n\| &= \|x_n + \theta_n(x_n - x_{n-1}) - x_n\| \\ &\leq \theta_n \|x_n - x_{n-1}\| \\ &= \theta_n \cdot \frac{\alpha_n}{\theta_n} \|x_n - x_{n-1}\| \rightarrow 0 \text{ as } n \rightarrow \infty, \end{aligned}$$

hence,

$$(3.19) \quad \lim_{i \rightarrow \infty} \|w_{n_i} - x_{n_i}\| = 0.$$

From (3.2) and (3.18), we have

$$\begin{aligned} \|y_n - w_n\| &= \|\beta_n w_n + (1 - \beta_n) T_{r_n}^{F,\phi} w_n - w_n\| \\ &\leq (1 - \beta_n) \|T_{r_n}^{F,\phi} w_n - w_n\| \rightarrow 0 \text{ as } n \rightarrow \infty, \end{aligned}$$

that is

$$(3.20) \quad \lim_{i \rightarrow \infty} \|y_{n_i} - w_{n_i}\| = 0.$$

It is easy to see from (3.19) and (3.20), that

$$(3.21) \quad \|y_{n_i} - x_{n_i}\| \leq \|y_{n_i} - w_{n_i}\| + \|w_{n_i} - x_{n_i}\| \rightarrow 0 \text{ as } i \rightarrow \infty.$$

Again from (3.2), we have

$$\begin{aligned} \|v_{n_i} - y_{n_i}\| &\leq \alpha \tau_n \|z_n\| \\ &= \alpha \frac{(\|y_n - S y_n\|^2 + \|A y_n - T A y_n\|^2)}{\|z_n\|}, \end{aligned}$$

thus by (3.12), we get

$$\|v_{n_i} - y_{n_i}\| \rightarrow 0 \text{ as } i \rightarrow \infty.$$

It follows from this and (3.19), that

$$(3.22) \quad \lim_{i \rightarrow \infty} \|v_{n_i} - w_{n_i}\| = 0.$$

By using (C1), (3.2) and (3.12), we derive

$$\begin{aligned} \|x_{n_i+1} - x_{n_i}\| &\leq \alpha_{n_i} \|x_{n_i} - f(x_{n_i})\| + (1 - \alpha_{n_i}) \alpha \tau_{n_i} \|z_{n_i}\| \\ &\leq \alpha_{n_i} \|x_{n_i} - f(x_{n_i})\| + (1 - \alpha_{n_i}) \alpha \frac{(\|y_{n_i} - S y_{n_i}\|^2 + \|A y_{n_i} - T A y_{n_i}\|^2)}{\|z_{n_i}\|^2}, \end{aligned}$$

which shows

$$(3.23) \quad \lim_{i \rightarrow \infty} \|x_{n_i+1} - x_{n_i}\| = 0.$$

We now show that  $\limsup_{i \rightarrow \infty} b_{n_i} \leq 0$ . Indeed, it suffices to show that

$$\limsup_{i \rightarrow \infty} \langle f(p) - p, x_{n_i+1} - p \rangle \leq 0.$$

Let  $\{x_{n_{i_j}}\}$  be a sequence of  $\{x_{n_i}\}$  such that

$$\lim_{j \rightarrow \infty} \langle f(p) - p, x_{n_{i_j}} - p \rangle = \limsup_{i \rightarrow \infty} \langle f(p) - p, x_{n_i} - p \rangle$$

Since  $\{x_{n_i}\}$  is bounded, there exists a subsequence  $\{x_{n_{i_j}}\}$  of  $\{x_{n_i}\}$  such that  $x_{n_{i_j}} \rightharpoonup q \in C$ . Without loss of generality, we may assume that  $x_{n_i} \rightharpoonup q$ , we obtain by (3.19), that  $w_{n_i} \rightharpoonup q$ . We

also have by (3.21) and (3.22) that  $y_{n_i}$  and  $v_{n_i}$  both converge weakly to  $q$ . Hence, by (3.14) and demiclosedness principle we have that  $q \in \text{Fix}(S)$ . Also, since  $A$  is a bounded linear operator we have that  $Ay_{n_i} \rightharpoonup Aq$ , thus by (3.14) again we obtain that  $Aq \in \text{Fix}(T)$ . Finally, we show that  $q \in \text{MEP}(F, \phi)$ . Let  $u_n = T_{r_n}^F w_n$ , we have by Lemma 2.3, that

$$F(u_n, y) + \phi(y) - \phi(u_n) + \frac{1}{r_n} \langle y - u_n, u_n - w_n \rangle \geq 0, \forall y \in H_1.$$

Now, since  $F$  is a monotone mapping, we obtain  $\phi(y) - \phi(u_n) + \frac{1}{r_n} \langle y - u_n, u_n - w_n \rangle \geq F(y, u_n)$  and hence  $\phi(y) - \phi(u_{n_i}) + \frac{1}{r_{n_i}} \langle y - u_{n_i}, u_{n_i} - w_{n_i} \rangle \geq F(y, u_{n_i})$  for all  $y \in H_1$ . It follows from (3.18), that  $u_{n_i} \rightharpoonup q$ . We obtain by (C3), (3.18) and the proper lower semicontinuity of  $\phi$  that

$$(3.24) \quad F(y, q) + \phi(q) - \phi(y) \leq 0, \forall y \in H_1.$$

Let  $y_t = ty + (1 - t)q$ , for all  $0 \leq t \leq 1$  and  $y \in H_1$ . It is easy to see that  $y_t \in H_1$ , thus (3.24) hold for  $y = y_t$ . that is

$$(3.25) \quad F(y_t, q) + \phi(q) - \phi(y_t) \leq 0.$$

From assumption (A1-A6) and (3.25), we have

$$\begin{aligned} 0 &= F(y_t, y) + \phi(y_t) - \phi(y) \\ &\leq tF(y_t, y) + (1 - t)F(y_t, q) + t\phi(y) + (1 - t)\phi(q) - t\phi(y_t) - (1 - t)\phi(y_t) \\ &= t[F(y_t, y) + \phi(y) - \phi(y_t)] + (1 - t)[F(y_t, q) + \phi(q) - \phi(y_t)] \\ &\leq t[F(y_t, y) + \phi(y) - \phi(y_t)]. \end{aligned}$$

Therefore, we obtain

$$(3.26) \quad t[F(y_t, y) + \phi(y) - \phi(y_t)] \geq 0, \forall y \in H_1.$$

Letting  $t \rightarrow 0$  in (3.26), obtain  $F(q, y) + \phi(y) - \phi(q) \geq 0, \forall y \in H_1$ , thus we have  $q \in \text{MEP}(F, \phi)$ . Hence  $q \in \Gamma$ .

From (3.8) and (3.23), we have

$$\begin{aligned} \limsup_{i \rightarrow \infty} \langle f(p) - p, x_{n_{i+1}} - p \rangle &\leq \limsup_{i \rightarrow \infty} \langle f(p) - p, x_{n_{i+1}} - x_{n_i} \rangle \\ &\quad + \limsup_{i \rightarrow \infty} \langle f(p) - p, x_{n_i} - p \rangle \\ &= \limsup_{j \rightarrow \infty} \langle f(p) - p, x_{n_j} - p \rangle \\ &= \langle f(p) - p, q - p \rangle \\ &\leq 0. \end{aligned}$$

We conclude by Lemma 2.5 that  $\{x_n\}$  converges strongly to a point  $p \in \Gamma$ , where  $p = P_\Gamma f(p)$ , The proof is complete.

#### 4. NUMERICAL EXAMPLE

In this section, we provide some numerical examples to illustrate the efficiency of Algorithm 1 and we compare the accelerated and non accelerated method for the SCFPPMEP.

**Example 4.1.** Let  $C = H_1 = \mathbb{R} = H_2$  and define the bounded linear operator  $A : H_1 \rightarrow H_2$  by  $Ax = 2x$  for all  $x \in \mathbb{R}$ . Define the bifunction  $F : C \times C \rightarrow \mathbb{R}$  by  $F(x, y) = 3x^2 + xy + 2y^2$

and  $\phi : C \rightarrow \mathbb{R}$  by  $\phi(x) = 0$ . Now, we compute  $u = T_r^{F,\phi}(x)$ . That is, we find  $u \in C$  such that for all  $z \in C$

$$\begin{aligned} 0 &\geq F_1(u, z) + \phi(u) + \frac{1}{r_n} \langle z - u, u - x \rangle \\ &= -3u^2 + uz + 2z^2 + \frac{1}{r_n} \langle z - u, u - x \rangle \end{aligned}$$

that is

$$\begin{aligned} 0 &\geq -3r_n u^2 + r_n uz + 2r_n z^2 + \langle z - u, u - x \rangle \\ &= -3r_n u^2 + r_n uz + 2r_n z^2 + uz - xz - u^2 + ux \\ &= 2r_n z^2 + (r_n u + u + z)z + (-3r_n u - u^2 + ux). \end{aligned}$$

Let  $h(z) = 2r_n z^2 + (r_n u + u - x)z + (-3r_n u^2 - u^2 + ux)$ . Then  $h(z)$  is a quadratic function of  $z$  with coefficients  $a = 2r_n$ ,  $b = r_n u + u - x$ , and  $c = -3r_n u^2 - u^2 + ux$ . We determine the discriminant  $\Delta$  of  $h(z)$  as follows:

$$\begin{aligned} \Delta &= (r_n u + u - x)^2 - 4(2r_n)(-3r_n u^2 - u^2 + ux), \\ &= 25r_n^2 u^2 + 10r_n u^2 + u^2 - 10r_n ux - 2ux + x^2, \\ (4.1) \quad &= ((5r_n + 1)u - x)^2. \end{aligned}$$

By Lemma 2.3,  $T_r^{F,\phi}$  is single-valued. Hence, it follows that  $h(z)$  has at most one solution in  $\mathbb{R}$ . Therefore, from (4.1) we have that  $u = \frac{x}{5r_n + 1}$ . This implies  $T_r^{F,\phi}(x) = \frac{x}{5r_n + 1}$  for all  $x \in H_1$ .

Define the mappings  $S : \mathbb{R} \rightarrow \mathbb{R}$  and  $T : \mathbb{R} \rightarrow \mathbb{R}$  by  $S(x) = -2x$  and  $T(x) = -3x$ , respectively. We set  $f(x) = \frac{x}{4}$ ,  $\beta_n = \frac{1}{2n+1}$ ,  $\alpha_n = \frac{1}{n+1}$ ,  $\epsilon_n = \frac{1}{3}$ ,  $\theta = \frac{1}{3}$ ,  $r_n = \frac{n+1}{2n}$  in Algorithm 1 for each  $n \in \mathbb{N}$ . It can easily be verified that all the condition of Theorem (3.2) are satisfied. We choose different initial values as follows:

Case 1  $x_0 = 1.78$ ,  $x_1 = 1.5$ ;

Case 2  $x_0 = 0.5$ ,  $x_1 = 0.15$ ;

Case 3  $x_0 = 0.05$ ,  $x_1 = 0.95$ .

## REFERENCES

- [1] H. A. ABASS, K. O. AREMU, L. O. JOLAOSO and O. T. MEWOMO, An inertial forward backward splitting method for solutions of certain optimization problems, *J. Nonlinear Funct. Anal.*, **2020** (2020), Article 6. [Online: <https://doi.org/10.23952/jnfa.2020.6>].
- [2] H. A. ABASS, C. IZUCHUKWU and O. T. MEWOMO, Viscosity approximation method for solutions of modified split generalized equilibrium, variational inequality and fixed point problems, *Rev. Un. Mat. Argentina*, **61** (2) (2020), pp. 389-411.
- [3] H.A. ABASS, C. IZUCHUKWU, O.T. MEWOMO and Q. L. DONG, Strong convergence of an inertial forward-backward splitting method for accretive operators in real Banach space, *Fixed Point Theory*, **21** (2) (2020), pp. 397-412.
- [4] H. A. ABASS and L. O. JOLAOSO, An inertial generalized viscosity approximation method for solving multiple-sets split feasibility problem and common fixed point of strictly pseudo-nonspreading mappings, *Axioms*, **10** (1) (2021). [Online: <https://doi.org/10.3390/axioms10010001>].




Figure 1: Example 4.1. Top left: Case 1, Top right Case 2, Bottom: Case 3.

- [5] H. A. ABASS, F. U. OGBUISI and O. T. MEWOMO, Common Solution of Split Equilibrium Problem and Fixed Point Problem With No Prior Knowledge of Operator Norm, *UPB Sci. Bull. A: Appl. Math. Phys.*, **80** (1) (2018), pp. 175-190.
- [6] T. O. ALAKOYA, L. O. JOLAOSO and O. T. MEWOMO, Modified inertial subgradient extragradient method with self adaptive stepsize for solving monotone variational inequality and fixed point problems, *Optimization*, (2020). [Online: [doi.org/10.1080/02331934.2020.1723586](https://doi.org/10.1080/02331934.2020.1723586)].
- [7] T. O. ALAKOYA, L. O. JOLAOSO and O. T. MEWOMO, Two modifications of the inertial Tseng extragradient method with self-adaptive step size for solving monotone variational inequality problems, *Demonstr. Math.*, (2020). [Online: [doi.org/10.1515/dema-2020-0013](https://doi.org/10.1515/dema-2020-0013)].
- [8] P. N. ANH and N. X. PHUONG, A parallel extragradient-like projection method for unrelated variational inequalities and fixed point problems, *J. Fixed Point Theory Appl.*, **20**, (2018), Article 74. [Online: <https://doi.org/10.1007/s11784-018-0554-1>].
- [9] Y. ARFAT, P. KUMAN, P. S. NGIASUMUNTHORN, M. A. A. KHAN, H. SARWAR and H. FUKHAR-UD-DIN, Approximation results for split equilibrium problems and fixed point problems of nonexpansive semigroup in Hilbert spaces, *Adv. in Differ. Equ.*, **512** (2020). [Online: <https://doi.org/10.1186/s13662-020-02956-8>].

- [10] E. BLUM and W. OETTLI, Optimization and variational inequalities to equilibrium problems, *Math. Stud.*, **63** (1994), pp. 123-145.
- [11] O. A. BOIKANYO, A Strongly Convergent Algorithm for the Split Common Fixed Point Problem, *Appl Math Comput.*, **265** (2015), pp. 844-853.
- [12] C. BYRNE, A Unified Treatment of Some Iterative Algorithms in Signal Processing and Image Reconstruction, *Inverse Probl.*, **20** (2004), pp. 103-120.
- [13] Y. CENSOR and A. SEGAL, The Split Common Fixed Point Problem for Directed Operators, *J. Convex Anal.*, **16** (2009), pp. 587-600.
- [14] X. CHEN, Y. SONG, J. HE and L. GONG, Self-Adaptive Algorithms for the Split Common Fixed Point Problem of the Demimetric Mappings, *Journal of Applied Mathematics and Physics*, **7** (2019), pp. 2187-2199
- [15] K. GOEBEL and S. REICH, *Uniform Convexity, Hyperbolic Geometry, Nonexpansive Mappings*, Marcel Dekker, New York, 1984.
- [16] Y. HAO, Some results of variational inclusion problems and fixed point problems with applications, *Appl. Math. Mech. (English Ed.)*, **30** (2009), pp. 1589-1596.
- [17] Z. HE, C. CHEN and F. GU, Viscosity approximation method for nonexpansive nonself-nonexpansive mappings and variational inequality, *J. Nonlinear Sci. Appl.*, **1** (2008), pp. 169-178.
- [18] H. IIDUKA and W. TAKAHASHI, Strong convergence theorems for nonexpansive mappings and inverse-strongly monotone mappings, *Nonlinear Anal.*, **61** (2005), pp. 341-350.
- [19] L. O. JOLAOSO, H. A. ABASS and O. T. MEWOMO, A Viscosity-Proximal Gradient Method with Inertial Extrapolation for Solving Minimization Problem in Hilbert Space, *Arch. Math. (Brno), Tomus*, **55** (2019), pp. 167-194.
- [20] I. V. KONNOV, S. SCHAIBLE and J. C. YAO, Combined relaxation method for mixed equilibrium problems, *J. Optim. Theory Appl.*, **126** (2005), pp. 309-322.
- [21] G. MARINO and H. K. XU, Weak and strong convergence theorems for strict pseudocontractions in Hilbert space, *J. Math. Anal. Appl.*, **329** (2007), pp. 336-346.
- [22] A. MOUDAFI, Viscosity approximation methods for fixed-point problems, *J. Math. Anal. Appl.*, **241** (2000), pp. 46-55.
- [23] A. MOUDAFI, Split monotone variational inclusions, *J. Optim. Theory Appl.*, **150** (2011), pp. 275-283.
- [24] A. MOUDAFI, A note on the split common fixed-point problem for quasi-nonexpansive operators, *Nonlinear Anal.*, **74** (2011), pp. 4083-4087.
- [25] O.K. OYEWOLE, H.A. ABASS and O.T. MEWOMO, A strong convergence algorithm for a fixed point constraint null point problem, *Rend. Circ. Mat. Palermo (2)*, (2020), [Online: <https://doi.org/10.1007/s12215-020-00505-6>].
- [26] B. T. POLYAK, Some methods of speeding up the convergence of iterative methods, *Zh. Vychisl. Mat. Mat. Fiz.*, **4** (1964), pp. 1-17.
- [27] W. PHUENGRATTANA and K. LERKCHAIYAPHUM, On solving the split generalized equilibrium problem and the fixed point problem for a countable family of nonexpansive multivalued mappings, *Fixed Point Theory Appl.* **2018** (2018), Article 6.[Online: <https://doi.org/10.1186/s13663-018-0631-6>].
- [28] M. RAHAMAN, Y. C. LIOU, R. AHMAD and I. AHMAD, Convergence theorems for split equality generalized mixed equilibrium problems for demi-contractive mappings, *J. Inequal. Appl.*, **2015** (2015), Article 418. [Online: <https://doi.org/10.1186/s13660-015-0936-5>].



- [29] S. SAEJUNG and P. YOTKAEW, Approximation of zeros of inverse strongly monotone operators in Banach spaces. *Nonlinear Anal.*, **75** (2012), pp. 724–750.
- [30] Y. SHEHU, Iterative Methods for Family of Strictly Pseudocontractive Mappings and System of Generalized Mixed Equilibrium Problems and Variational Inequality Problems, *Fixed Point Theory Appl.*, **22** (2011). [Online: <https://doi.org/10.1155/2011/852789>].
- [31] S. SUANTAI, P. JAILOKA and A. HANJING, An accelerated viscosity forward-backward splitting algorithm with the linesearch process for convex minimization problems, *J. Inequal. Appl.*, **42** (2021). [Online: <https://doi.org/10.1186/s13660-021-02571-5>].
- [32] W. TAKAHASHI, *Nonlinear functional analysis*, Yokohama Publishers, Yokohama, 2000.
- [33] W. TAKAHASHI and M. TOYODA, Weak convergence theorems for nonexpansive mappings and monotone Mappings, *J. Optim. Theory Appl.*, **118** (2003), pp. 417-428.
- [34] F. WANG, A New Iterative Method for the Split Common Fixed Point Problem in Hilbert Spaces. *Optimization*, **66** (2017), pp. 407-415.
- [35] F. WANG and H. K. XU, Cyclic Algorithms for Split Feasibility Problems in Hilbert Spaces, *Nonlinear Anal.*, **74** (12) (2011) pp. 4105-4111.
- [36] Y. H. YAO, J. C. YAO, Y. C. LIOU and M. POSTALACHE, Iterative Algorithms for Split Common Fixed Points of Demicontractive Operators without Prior Knowledge of Operator Norms, *Carpathian J. Math.*, **34** (2018), pp. 459-466.
- [37] Y. H. YAO, M. POSTALACHE, Y. C. LIOU and Z. S. YAO, Construction Algorithms for a Class of Monotone Variational Inequalities, *Optim. Lett.*, **10** (2016), pp. 1519-1528.